

I ILLINOIS


Veterinary Teaching Hospital
COLLEGE OF VETERINARY MEDICINE

Transdermal Medication Administration Tips

- Wear gloves when applying the medication
- Apply the prescribed amount of medication to the pinna of one ear
- Rub the medication in completely
- Alternate ears with each dose
- Clean the ear prior to applying the next dose (use water or a generic ear cleaning solution)
- Irritation is the most common adverse effect with any transdermal. Cleaning the ear prior to each dose and alternating ears is the most effective way to prevent irritation.
- Keep small children and other pets away from the application site for at least 2 hours after each dose.

Small Animal Clinic | 217-333-5300
1008 W. Hazelwood Dr., Urbana

Large Animal Clinic | 217-333-2000
1102 W. Hazelwood Dr., Urbana



I ILLINOIS

Veterinary Teaching Hospital
COLLEGE OF VETERINARY MEDICINE

Transdermal Medication Administration Tips

- Wear gloves when applying the medication
- Apply the prescribed amount of medication to the pinna of one ear
- Rub the medication in completely
- Alternate ears with each dose
- Clean the ear prior to applying the next dose (use water or a generic ear cleaning solution)
- Irritation is the most common adverse effect with any transdermal. Cleaning the ear prior to each dose and alternating ears is the most effective way to prevent irritation.
- Keep small children and other pets away from the application site for at least 2 hours after each dose.

Small Animal Clinic | 217-333-5300
1008 W. Hazelwood Dr., Urbana

Large Animal Clinic | 217-333-2000
1102 W. Hazelwood Dr., Urbana

